

Tips for Patients Using Insulin Pumps

Experts agree with the following steps*:

- * Choose a part of the body
- * Divide into 4 areas
- * Use one area every 7-10 days depending on your pump set type
- * Always use a new site
- * Avoid using a single site more than once every 4 weeks
- * Follow your health care professional's recommendations for site rotation




Use for diabetes-related injection-induced scarring

Scarring from on-going use of wearables can significantly impact glycemic control

- ▶ GREATER glycemic variability up to 7 times higher*
- ▶ INCREASED use of insulin up to 35% more[†]


▶ HIGHER A1c level - up to 0.55 higher*



STEP 1: Apply

Holding embrace* on its outer edges, use the Peel Tabs to peel off the clear protective liner that covers the adhesive patch.

DO NOT touch the adhesive patch with your fingers.



Position embrace® over the injection/scarred area so that the Applicator is parallel to the length of the area and faces away from your skin.

embrace® against the skin for 60 seconds. Make sure to press down firmly on the edges and corners.



STEP 2: Release

Use one hand to hold embrace® in place. Use the other hand to press firmly across the entire surface of the Applicator to ensure the patch is adhered well to the skin.



Holding embrace® against your skin with one hand, use your other hand to pull one Release Tab to the side and along the edge of the Applicator.

Repeat this step with the other Release Tab.
The Applicator will completely detach, leaving the patch adhered to the skin.

Smooth out any uneven areas, flatten any air pockets, and secure any raised corners or edges.



Example for Omnipod site rotation with embrace medium




Area = 3 pod sites Insert 3 pods aligned in area of choice



After removing third pod, immediately apply embrace over injured sites to PROTECT from scarring. Wear 9 days while pumping in next area



Rotate pods to new area and repeat treatment with embrace at previously used site




Continue rotation pattern with each new area used

Example for 3 Day pump infusion set site rotation with embrace medium



Area = 3 pod sites
Insert 3 pods aligned in area
of choice



After removing third pod, immediately apply embrace over injured sites to PROTECT from scarring.

Wear 9 days while pumping in next area




Rotate pods to new area and repeat treatment with embrace at previously used site



Continue rotation pattern with each new area used


Example for 7 Day pump infusion set site rotation with embrace small




Insert pumpset in area of choice



After removing, immediately apply embrace over injured site to PROTECT from scarring. Wear 7 days while pumping in next area



Rotate pumpset to new area and repeat treatment with embrace at previously used site



Continue rotation pattern with each new area used

We're here to support your use of embrace® to make sure you experience the best possible results. Contact us for questions or additional information.



^{*} Frid, A. H., Hirsch, L. J., Menchior, A. R., Morel, D. R. & Strauss, K.W. Worldwide Injection Technique Questionnaire Study: Injecting Complications and the Role of the Professional. Mayo Clin Proc 91, 1224-1230, doi:10.1016/j.mayocp.2016.06.012 (2016).
† Blanco M, Hernández MT, Strauss KW, Amaya M. Prevalence and risk factors of lipohypertrophy in insulin-injecting patients with diabetes. Diabetes Metab. 2013 Oct;39(5):445-53. doi: 10.1016/j.diabet.2013.05.006. Epub 2013 Jul 22. PMID: 23886784.

^{*} Frid AH, Kreugel G, Grassi G, et.al. New insulin delivery recommendations. Mayo Clic. Proc. 2016-91(9): 1231-1255